



# Reumatología Clínica



<https://www.reumatologiaclinica.org>

## P019 - MAINTENANCE OF CLINICAL RESPONSE IN INDIVIDUAL CHILDREN WITH JUVENILE IDIOPATHIC ARTHRITIS TREATED WITH SUBCUTANEOUS ABATACEPT

I. Calvo Penadés<sup>1</sup>, N. Ruperto<sup>2</sup>, H. Brunner<sup>3</sup>, N. Tzaribachev<sup>2</sup>, I. Louw<sup>2</sup>, F. Zapata<sup>4</sup>, G. Horneff<sup>2</sup>, I. Foeldvari<sup>2</sup>, D. Kingsbury<sup>3</sup>, R. Joos<sup>2</sup>, M.E. Paz Gastanaga<sup>2</sup>, C. Wouters<sup>2</sup>, J. Breedt<sup>2</sup>, T. Lutz<sup>2</sup>, T. Miraval<sup>2</sup>, N. Rubio<sup>2</sup>, Y. Elbez<sup>5</sup>, M. Nys<sup>6</sup>, R. Wong<sup>7</sup>, A. Martini<sup>2</sup> and D.J. Lovell<sup>3</sup> <sup>poi</sup>

<sup>1</sup>Hospital Universitario y Politécnico La Fe. Valencia. <sup>2</sup>PRINTO. Istituto Gaslini. Genoa (Italy). <sup>3</sup>PRCSG. CHMC Cincinnati (USA). <sup>4</sup>PRINTO. Star Medica Hospital. Mérida Yucatán (México). <sup>5</sup>Excelya. Boulogne-Billancourt (France). <sup>6</sup>Bristol-Myers Squibb. Braine-L'Alleud (Belgium). <sup>7</sup>Bristol-Myers Squibb. Princeton. NJ (USA).

### Resumen

**Introduction:** The efficacy of SC abatacept (ABA) in patients (pts) with polyarticular-course juvenile idiopathic arthritis (pJIA) was shown in a 2-year (yr), Phase III, open-label international study (NCT01844518). However, it is unknown whether each individual pt within a treatment group consistently achieves the same efficacy endpoint at all time points.

**Objectives:** To investigate whether ABA efficacy is maintained by individual pts with pJIA over time.

**Methods:** In this subgroup analysis, pts in two age cohorts (2-5 yrs and 6-17 yrs) who achieved clinical response to weekly SC ABA (10 to 25 kg [50 mg], 25 to 50 kg [87.5 mg], ? 50 kg [125 mg]) at Day 113 (time point of primary pharmacokinetics endpoint<sup>1</sup>) were followed for 2 yrs. Stringent efficacy outcomes selected for analysis included JIA-ACR70, JIA-ACR100, Juvenile Arthritis Disease Activity Score 71 (JADAS71) minimal disease activity (MDA; ? 3.8) and JADAS71 inactive disease (ID; ? 1) rates. Prognostic factors for sustained response were investigated using logistic regression.

**Results:** A total of 219 pts entered the study (46 [21.0%] 2-5 yrs; 173 [79.0%] 6-17 yrs) and a subgroup of these pts achieved a clinical response at Day 113 (Table 1). Most pts who achieved JIA-ACR70, JIA-ACR100, JADAS71 MDA and JADAS71 ID at Day 113 sustained their response at Day 393 and at Days 393 and 645 in the 2-5-yr and 6-17-yr cohorts (Table 2). Across cohorts, more than 75% and 60% of pts maintained a JIA-ACR 70 and JADAS71 MDA response through Day 645, respectively. Prior biologic (b)DMARD use was an important prognostic factor. In pts aged 6-17 yrs, sustained JIA-ACR70 response rate at Days 393 and 645 was 81% (57/70) in pts who did not have prior bDMARDs vs 57% (12/21) in pts who had prior bDMARDs (p = 0.0715); sustained JADAS71 MDA response rate was 71% (37/52) vs 37% (7/19; p = 0.0320). Prognostic factors for JIA-ACR100 response and JADAS71 ID in pts aged 6-17 yrs and for all outcomes in pts aged 2-5 yrs could not be determined due to low pt numbers.

Table 1. Proportion of pts who achieved a clinical response at Day 113

Endpoint	Pts with response at Day 113	
	2-5 yrs (n = 46)	6-17 y (n = 173)
JIA-ACR70	34 (74)	91 (53)
JIA-ACR100	19 (41)	25 (15)
JADAS71 MDA	28 (61)	71 (41)
JADAS71 ID	17 (37)	28 (16)

Data are expressed as n (%). ID: inactive disease; JADAS71: Juvenile Arthritis Disease Activity Score 71; MDA: minimal disease activity; pt: patient; yr: year.

Table 2. Proportion of responders at Day 113 with sustained clinical response at Day 393, and at Days 393 and 645 in cohort aged 2-5 yrs and cohort aged 6-17 yrs

	2-5-yr-old pts who achieved a clinical response at Day 113				6-17-yr-old pts who achieved a clinical response at Day 113			
	JIA-ACR70 (n = 34)	JIA-ACR100 (n = 19)	JADAS71 MDA (n = 28)	JADAS71 ID (n = 17)	JIA-ACR70 (n = 91)	JIA-ACR100 (n = 25)	JADAS71 MDA (n = 71)	JADAS71 ID (n = 28)
Day 393	32 (94)	12 (63)	25 (89)	10 (59)	78 (86)	20 (80)	58 (82)	21 (75)
Day 645	32 (94)	11 (58)	23 (82)	9 (53)	69 (76)	15 (60)	44 (62)	18 (64)

Data are expressed as n (%). ID: inactive disease; JADAS71: Juvenile Arteritis Disease Activity Score 71; MDA: minimal disease activity; pt: patient; yr: year.

**Conclusions:** The majority of individuals with pJIA who achieved stringent efficacy endpoints with weekly SC abatacept by Day 113 sustained that clinical endpoint over time. Prior bDMARD use may be a prognostic factor for sustained response over 2 yrs.

Código EUDRACT: NCT01844518.

## References

1. Brunner HI, et al. Arthritis Rheumatol. 2018;70:1144-54. Abstract presented at EULAR 2019. Published in Ann Rheum Dis. 2019. DOI: 10.1136/annrheumdis-2019-eular.236.