



Sociedad Española
de Reumatología -
Colegio Mexicano
de Reumatología

Reumatología Clínica

www.reumatologiaclinica.org



Original Article

Association between serum dickkopf-1 levels and disease duration in axial spondyloarthritis



Roxana Rubio Vargas^{a,*}, Enrique Melguizo Madrid^b, Concepción González Rodríguez^b, Federico Navarro Sarabia^a, Carmen Dominguez Quesada^a, Rafael Ariza Ariza^a, Victoria Navarro Compán^c

^a Department of Rheumatology, University Hospital Virgen Macarena, Seville, Spain

^b Biochemistry Department, University Hospital Virgen Macarena, Seville, Spain

^c Department of Rheumatology, IdiPaz, University Hospital La Paz, Madrid, Spain

ARTICLE INFO

Article history:

Received 26 January 2016

Accepted 29 April 2016

Available online 11 June 2016

Keywords:

DKK-1

Disease duration

Axial spondyloarthritis

ABSTRACT

Background: Axial spondyloarthritis (axSpA) is characterized by new bone formation. The complex systems underlying this process involve Wnt-signaling pathway. It has been observed that serum levels of dickkopf-1 (DKK-1), an important inhibitor of Wnt-signaling, are decreased in patients with axSpA. However, these data are from studies including only patients with long-standing disease.

The aim of this study is to investigate if symptom duration influences on serum DKK-1 levels in patients with axSpA.

Material and methods: A cross-sectional study including consecutive patients with axSpA (ASAS criteria) naïve for anti-TNF therapy. Collected data included demographic and disease characteristics, time since first symptom onset, assessment of disease activity and function, and determination of DKK-1 serum levels. Patients were classified as early axSpA (symptom duration ≤ 5 years) and established axSpA (>5 years). Linear regression models were employed to investigate the variables related to DKK-1 serum levels.

Results: In total, 90 patients were included. Sixty-eight patients had early axSpA and 22 had established disease. Serum levels of DKK-1 were significantly higher in patients with early axSpA compared with established axSpA (22.1 ± 12.6 vs 16.4 ± 10.7 pM; $p = 0.04$). Among all tested variables, only symptom duration was significantly and inversely correlated with DKK-1 serum levels ($\beta = -0.041$; $p = 0.01$).

Conclusion: Serum DKK-1 levels in axSpA depend on disease duration. As disease duration increases, DKK-1 serum levels decrease. Based on this, an intensive treatment at early stages of the disease could have a better outcome on inhibiting/slowng radiographic progression in patients with axSpA.

© 2016 Elsevier España, S.L.U. and Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. All rights reserved.

Asociación entre niveles séricos de Dickkopf-1 y duración de la enfermedad en espondiloartritis axial

RESUMEN

Palabras clave:

DKK-1

Duración de la enfermedad

Espondiloartritis axial

Objetivo: Espondiloartritis axial (EsPax) se caracteriza por nueva formación ósea. El complejo sistema que subyace este proceso incluye la vía de señal Wnt. Se ha demostrado que niveles séricos de dickkopf-1 (DKK-1), un importante inhibidor de la vía de señal Wnt, está disminuidos en pacientes con EsPax. Sin embargo, estos datos proceden exclusivamente de pacientes con EsPax de larga duración.

El objetivo de este estudio es investigar si la duración de la enfermedad influye en niveles séricos de DKK-1 en pacientes con EsPax.

Material y métodos: Estudio transversal en pacientes con EsPax sin terapia anti-TNF. Se recogieron datos demográficos y de la enfermedad, y se determinaron niveles de DKK-1 séricos en la misma visita. Los

* Corresponding author.

E-mail address: roxanarubio27@gmail.com (R. Rubio Vargas).

pacientes fueron clasificados en base a la duración de síntomas en EsPax precoz (≤ 5 años) y establecida (> 5 años). Se emplearon modelos de regresión lineal para investigar las variables asociadas con los niveles de DKK-1.

Resultados: Se incluyeron 90 pacientes, 68 con EsPax precoz y 22 con EsPax establecida. Los niveles de DKK-1 fueron superiores en EsPax precoz comparado con EsPax establecida (22.1 ± 12.6 vs 16.4 ± 10.7 pM; $p = 0.04$). De todas las variables, sólo la duración de síntomas se asoció significativamente con DKK-1 (beta: -0.041 ; $p = 0.01$).

Conclusiones: Los niveles séricos de DKK-1 en EsPax, dependen de la duración de la enfermedad. A medida que la duración de la enfermedad aumenta, niveles séricos de DKK-1 disminuye. Por lo tanto, el tratamiento intensivo en estadios tempranos de la enfermedad podría tener un mejor resultado en inhibir/disminuir la progresión radiológica en pacientes con EsPax.

© 2016 Elsevier España, S.L.U.

y Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. Todos los derechos reservados.

Introduction

Axial spondyloarthritis (axSpA) is a chronic inflammatory disease that mainly affects the spine and sacroiliac joints. Structural damage in axSpA is characterized by new bone formation leading to syndesmophytes, bone bridges and complete ankylosis. The complex systems that underlie new bone formation in patients with axSpA involve Wingless protein (Wnt) signaling and growth factors such as the bone morphogenetic proteins (BMPs).

An important inhibitor of the Wnt signaling is dickkopf-related protein 1 (DKK-1), which seems to be enhanced by tumor necrosis factor alpha (TNF). Serum DKK-1 levels are decreased in patients with axSpA compared with healthy controls¹ and other rheumatic inflammatory diseases such as rheumatoid arthritis.² Additionally, it has been shown that DKK-1 serum levels are related to radiographic progression in patients with axSpA.³ Specifically, high levels of functional DKK-1 seems to be protective of the development of syndesmophytes,⁴ and DKK-1 blockade leads to the fusion of sacroiliac joints in an animal model of arthritis.⁵

Furthermore, the mechanisms and characteristics influencing serum DKK-1 levels in patients with axSpA are unclear. In principle, the most accepted hypothesis is that TNF induces DKK-1,⁶ which down-regulates bone formation through inhibition of Wnt signaling. However, the change produced in DKK-1 serum levels after treatment with TNF blockers is not yet conclusive since different results (decrease, increase and even no change on DKK-1) have been reported in this sense.^{7–9} In addition, TNF blockers have shown clinical efficacy in patients with axSpA,¹⁰ but inhibition or slowing radiographic progression with this therapy has not been demonstrated.^{11–13} Based on this, it may be hypothesized that inflammation and radiographic progression are uncoupled processes in axSpA.

Furthermore, all these data come from studies including patients with long-standing and established disease.^{11–13} The influence of disease duration on DKK-1 levels is unknown since no study has previously assessed whether or not patients at early stages of the disease have the same decreased levels of DKK-1 than patients with longstanding and established disease. Based on this, the aim of this study is to investigate if disease duration influences on serum levels of DKK-1 in patients with axSpA.

Materials and methods

Study design, population and data collection

A cross-sectional study in which consecutive patients with axSpA attending the outpatient clinic of a tertiary hospital between January 2011 and June 2014 were included. Patients should fulfill Assessment of SpondyloArthritis International Society (ASAS)

classification criteria for axSpA and be naïve for treatment with TNF blockers or other biological therapy.

The following data were collected: demographic characteristics including age and gender; disease characteristics specifying symptoms duration, human leukocyte antigen B27 (HLA-B27) and presence or previous history of SpA features including psoriasis, inflammatory bowel disease, uveitis or peripheral arthritis; and concomitant treatment. Disease Activity was measured by the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI), the Ankylosing Spondylitis Disease Activity Score (ASDAS), patient's visual analog score of disease activity (Patient's VAS) and C-Reactive Protein (CRP). Enthesitis was assessed using the Maastricht Ankylosing Spondylitis Enthesitis Score (MASES) and function was evaluated through the Bath Ankylosing Spondylitis Functional Index (BASFI). A physical examination was also performed to assess mobility, measured by the Bath Ankylosing Spondylitis Mobility Index (BASMI). Blood samples to determine Dkk-1 serum levels by enzyme immunoassay were collected too. In addition, based on symptoms duration patients were classified in two groups: early axSpA (less than or equal to 5 years) and established axSpA (more than 5 years). This study was approved by the local ethical committee, and all patients signed a written informed consent before being included.

Quantification of dickkopf-1 (Dkk-1) levels

Serum samples were obtained through venous punctures and stored at -80°C until assays were performed. Commercially available kit was used (Biomedica Medizinprodukte GmbH & Co KG, A-1210 Wien, Divischgasse 4). Serum samples were diluted 2:7 and were processed by an enzyme immunoassay. DKK-1 levels were estimated by means of colourimetric measurements at 450 nm and by interpolation from a standard curve (0–160 pM) on an ELISA plate reader, obtaining DKK-1 serum concentrations in pM. The intra- and inter-assay coefficients of variation were $< 3\%$.

Statistical analysis

First, a descriptive analysis was performed. For this purpose, collected variables between patients with early and established axSpA were compared using Student *t*-test for continuous variables and chi-square test for categorical variables. The results are presented as mean and standard deviation (SD) for continuous variables and as percentage and relative frequencies for categorical variables. Second, separate linear regression models adjusted for age and gender were employed for identifying the variables related to serum levels of DKK-1. For all the analyses, SPSS software version 21.0 was

Table 1
Demographic, disease characteristics and biomarkers based on symptoms duration.

	Early axSpA N = 68	Established axSpA N = 22	p value
Age (years)	34.6 ± 9.7	43.2 ± 11	<0.01
Gender (male, %)	48 (70.6%)	20 (90.9%)	<0.05
Symptoms duration (months)	19.3 ± 17.4	183.3 ± 112.4	<0.001
Uveitis	3 (4.4%)	7 (31.8%)	<0.01
Peripheral arthritis	38 (55.9%)	14 (63.6%)	0.5
Psoriasis	4 (5.9%)	4 (18.2%)	0.09
Inflammatory bowel disease	9 (14.8%)	1 (4.8%)	0.2
HLA-B27	44 (68.8%)	17 (85%)	0.1
BASDAI (0–10)	5.3 ± 2.3	7.0 ± 1.8	0.001
ASDAS-CRP	2.9 ± 1.2	3.8 ± 0.8	<0.001
CRP (mg/L)	10.5 ± 17.2	17.8 ± 21.5	0.1
Patient's VAS (0–100)	50.5 ± 27.7	70.0 ± 22.9	<0.01
BASFI (0–10)	4.1 ± 2.9	6.5 ± 2.4	<0.001
MASES (0–13)	1.4 ± 2.6	3.5 ± 2.9	<0.01
BASMI (0–10)	2.7 ± 1.3	4.1 ± 1.4	<0.001
Concomitant NSAID	40 (58.8%)	15 (68.2%)	<0.01
DKK-1 (pM)	22.1 ± 12.6	16.4 ± 10.7	0.04

BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; ASDAS: Ankylosing Spondylitis Disease Activity Score; CRP: C Reactive Protein; Patient's VAS: patient's global visual analog scale; BASFI: Bath Ankylosing Spondylitis Functional Index; MASES: Maastricht Ankylosing Spondylitis Enthesitis Score; BASMI: Bath Ankylosing Spondylitis Metrology Index; NSAID: Nonsteroidal Anti-Inflammatory Drugs; DKK-1: Dickkopf-1. Results are shown as mean ± standard deviation or absolute number with percentages. p value in bold: with statistical significance.

employed and p-values less than 0.05 were considered statistically significant.

Results

Ninety patients with axSpA were included in this study. Sixty-eight patients were classified as early axSpA and 22 as established disease. Results for demographic, disease characteristics and biomarkers in each group of patients are shown in Table 1. Compared to patients with established disease, patients with early axSpA were younger, had a higher proportion of females, lower degree of disease activity and less enthesitis involvement. As expected, they also had higher mobility range aside from a better function.

Importantly, DKK-1 serum levels were found to be statistically significant higher in patients with early axSpA compared with patients with established disease (22.1 ± 12.6 vs 16.4 ± 10.7 pM; $p = 0.04$, respectively).

Furthermore, Table 2 shows the results for the separate linear regression models evaluating the association between DKK-1 serum levels and symptom duration, disease activity, function and mobility. Among all the tested variables, only symptom duration was statistically significant related to DKK-1 serum levels (standard β : -0.041; $p = 0.01$). This relationship was inverse: as disease duration increased, DKK-1 serum levels decreased.

Discussion

This study shows that DKK-1 serum levels in patients with axSpA depend on symptom duration. Independently of the age and gender, patients with early axSpA have higher serum DKK-1 levels compared with patients with established disease.

These findings are consistent with published data, in which low serum levels of DKK-1 in patients with established ankylosing spondylitis have been observed.^{1,14,15} On the other hand, opposite to a previous study, an association between disease activity and DKK-1 levels was not found.¹⁵ In our study, most patients had active or very active disease activity while in the study of Korskos et al. the degree of disease activity was more heterogeneous.

Table 2
Separate linear regression models for the association between DKK-1 and disease characteristics.

	Beta	Std. error	p value
Symptom duration (months)	-0.041	0.016	<0.01
HLA-B27 positive	-2.626	3.129	0.4
Psoriasis	-1.281	4.671	0.8
Inflammatory bowel disease	-3.214	4.386	0.5
Uveitis	2.454	4.253	0.6
Peripheral arthritis	0.809	2.687	0.8
BASDAI (0–10)	-0.285	0.589	0.6
ASDAS (CRP)	0.034	1.231	1.0
CRP (mg/L)	0.055	0.074	0.5
Patient's VAS (0–100)	-0.014	0.050	0.8
BASFI (0–10)	0.075	0.479	0.9
MASES (0–13)	0.100	0.499	0.8
BASMI (0–10)	0.478	0.041	0.6

BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; ASDAS: Ankylosing Spondylitis Disease Activity Score; CRP: C Reactive Protein; Patient's VAS: patient's global visual analog scale of disease activity; BASFI: Bath Ankylosing Spondylitis Functional Index; MASES: Maastricht Ankylosing Spondylitis Enthesitis Score; BASMI: Bath Ankylosing Spondylitis Metrology Index; Beta: unstandardized beta coefficient; Std. error: standard error. All models are adjusted for age and gender. p value in bold: with statistical significance.

This may explain the difference on the results between the studies.

The difference on DKK-1 serum levels found in our study between patients with early and established axSpA may be relevant to understand why despite suppressing inflammation, TNF blockers have not shown to inhibit the development of syndesmophytes in patients with axSpA. On one hand, disease activity has been longitudinally related to radiographic progression in patients with axSpA.¹⁶ On the other hand, TNF blockers decrease disease activity but have not shown to inhibit new bone formation in patients with axSpA.^{11–13}

In addition, it has been shown that low levels of serum DKK-1 are necessary to develop syndesmophytes.⁴ What is unclear is the relationship between TNF and DKK-1. TNF stimulates DKK-1 but previous studies have found that inhibition of TNF by TNF-blockers does not necessarily lead to the decrease of serum DKK-1.^{7,9} This highlights the fact that there may be more important factors different than TNF regulating DKK-1 production. As shown in the results of our study, in patients with established axSpA DKK-1 serum levels are already decreased.¹⁷ This may explain why despite suppressing disease activity, the administration of a more intensive therapy such as TNF-blockers in a late stage of the disease is not able to avoid the development of syndesmophytes. Nevertheless, this situation may be different in patients at an early stage of the disease, in which DKK-1 serum levels are not decreased yet. Based on this, initiation of TNF-blockers at early stages of the disease in patients with axSpA could have a different (better) outcome on inhibiting or slowing radiographic progression. So, these results support the hypothesis of a window of opportunity to treat patients with axSpA if the avoidance of irreversible radiographic damage is desired.

The following process has been postulated to explain how inflammation could lead to new bone formation in axSpA¹⁸: First, the initial development of inflammation causes erosive cartilage and bone destruction. Later, these lesions are filled in by fibrous tissue. And finally, this tissue is ossified leading to syndesmophytes. This process is mediated by TNF- α and Dkk-1. TNF-blockers target the sites of active inflammation but have no direct effects on osteoproliferation.¹⁸ In other words, it could be postulated that TNF-blockers block the transformation from inflammation to erosion; however, they don't block the transformation from erosion to osteoproliferation, where non-steroidal anti-inflammatory drugs might have a role.¹⁹ This may explain why initiation of

TNF-blockers at early stages of the disease could have a better outcome on the development of radiographic progression.

Furthermore, several limitations of this study should be taken into account when interpreting its results. First, this is a cross-sectional study, which does not allow investigating if DKK-1 serum levels truly decrease overtime. Second, no control group was included so it is not possible to know whether patients with early axSpA have similar or lower serum levels of DKK-1 than healthy population. Third, the group of patients with established disease was substantially smaller than the group with early disease and a bigger sample size in this group could have an impact. Finally, no radiographs were available to assess radiographic damage or progression in these patients.

In summary, DKK-1 serum levels depend on symptom duration in patients with axSpA. Patients with early disease have higher serum DKK-1 levels compared with patients with established axSpA. Based on the results of the current study, it can be concluded that serum DKK-1 levels are not decreased from the beginning of the disease in patients with axSpA but they decrease as disease duration increases. This could underlie a window of opportunity to avoid development of new bone formation through TNF-blockers in patients with axSpA. Nevertheless, further data from longitudinal studies are necessary to confirm this.

Ethical responsibilities

Protection of people and animals. The authors state that the procedures conformed to the ethical standards of human experimentation committee responsible and according to the World Medical Association and the Declaration of Helsinki.

Confidentiality of data. The authors declare that they have followed the protocols of the workplace on the publication of patient data.

Right to privacy and informed consent. The authors declare that this article does not appear patient data.

Funding

This work was supported by an Initiated Investigator Research (IIR) grant from Pfizer, the Institute Carlos III (PI10/01963) and the Spanish Society of Rheumatology (call 2010).

Conflicts of interest

All authors declare no conflicts of interest in this manuscript.

Acknowledgments

We are grateful to all the patients that participated in this study for their collaboration. In addition, we would like to thank Cristian

Leyva, Rufino Mondéjar, Virginia Moreira and Consuelo Leon, who participated in the determination and collection of clinical and laboratory variables. We also thank Ismael Gómez for medical advisory support.

References

1. Kwon S-R, Lim M-J, Suh C-H, Park S-G, Hong Y-S, Yoon B-Y, et al. Dickkopf-1 level is lower in patients with ankylosing spondylitis than in healthy people and is not influenced by anti-tumor necrosis factor therapy. *Rheumatol Int.* 2012;32:2523–7.
2. Wang S-Y, Liu Y-Y, Ye H, Guo J-P, Li R, Liu X, et al. Circulating Dickkopf-1 is correlated with bone erosion and inflammation in rheumatoid arthritis. *J Rheumatol.* 2011;38:821–7.
3. Yucong Z, Lu L, Shengfa L, Yongliang Y, Ruguo S, Yikai L. Serum functional dickkopf-1 levels are inversely correlated with radiographic severity of ankylosing spondylitis. *Clin Lab.* 2014;60:1527–31.
4. Heiland GR, Appel H, Poddubnyy D, Zwerina J, Hueber A, Haibel H, et al. High level of functional dickkopf-1 predicts protection from syndesmophyte formation in patients with ankylosing spondylitis. *Ann Rheum Dis.* 2012;71:572–4.
5. Uderhardt S, Diarra D, Katzenbeisser J, David J-P, Zwerina J, Richards W, et al. Blockade of Dickkopf(DKK)-1 induces fusion of sacroiliac joints. *Ann Rheum Dis.* 2010;69:592–7.
6. Wendling D, Claudepierre P. New bone formation in axial spondyloarthritis. *Jt Bone Spine Rev Rhum.* 2013;80:454–8.
7. Daoussis D, Lioussis S-NC, Solomou EE, Tسانaktsi A, Bounia K, Karampetsou M, et al. Evidence that Dkk-1 is dysfunctional in ankylosing spondylitis. *Arthritis Rheum.* 2010;62:150–8.
8. Korkosz M, Gąsowski J, Leszczyński P, Pawlak-Buś K, Jeka S, Siedlar M, et al. Effect of tumour necrosis factor- α inhibitor on serum level of dickkopf-1 protein and bone morphogenetic protein-7 in ankylosing spondylitis patients with high disease activity. *Scand J Rheumatol.* 2014;43:43–8.
9. Taylan A, Sari I, Akinci B, Bilge S, Kozaci D, Akar S, et al. Biomarkers and cytokines of bone turnover: extensive evaluation in a cohort of patients with ankylosing spondylitis. *BMC Musculoskelet Disord.* 2012;13:191.
10. Callhoff J, Sieper J, Weiß A, Zink A, Listing J. Efficacy of TNF α blockers in patients with ankylosing spondylitis and non-radiographic axial spondyloarthritis: a meta-analysis. *Ann Rheum Dis.* 2015;74:1241–8.
11. van der Heijde D, Landewé R, Baraliakos X, Houben H, van Tubergen A, Williamson P, et al. Radiographic findings following two years of infliximab therapy in patients with ankylosing spondylitis. *Arthritis Rheum.* 2008;58:3063–70.
12. van der Heijde D, Landewé R, Einstein S, Ory P, Vosse D, Ni L, et al. Radiographic progression of ankylosing spondylitis after up to two years of treatment with etanercept. *Arthritis Rheum.* 2008;58:1324–31.
13. van der Heijde D, Salonen D, Weissman BN, Landewé R, Maksymowych WP, Kupper H, et al. Assessment of radiographic progression in the spines of patients with ankylosing spondylitis treated with adalimumab for up to 2 years. *Arthritis Res Ther.* 2009;11:R127.
14. Diarra D, Stolina M, Polzer K, Zwerina J, Ominsky MS, Dwyer D, et al. Dickkopf-1 is a master regulator of joint remodeling. *Nat Med.* 2007;13:156–63.
15. Korkosz M, Gąsowski J, Leszczyński P, Pawlak-Buś K, Jeka S, Kucharska E, et al. High disease activity in ankylosing spondylitis is associated with increased serum sclerostin level and decreased wingless protein-3a signaling but is not linked with greater structural damage. *BMC Musculoskelet Disord.* 2013;14:99.
16. Ramiro S, van der Heijde D, van Tubergen A, Stolwijk C, Dougados M, van den Bosch F, et al. Higher disease activity leads to more structural damage in the spine in ankylosing spondylitis: 12-year longitudinal data from the OASIS cohort. *Ann Rheum Dis.* 2014;73:1455–61.
17. Ustun N, Tok F, Kalyoncu U, Motor S, Yuksel R, Yagiz AE, et al. Sclerostin and Dkk-1 in patients with ankylosing spondylitis. *Acta Reumatol Port.* 2014;39:146–51.
18. Sieper J, Appel H, Braun J, Rudwaleit M. Critical appraisal of assessment of structural damage in ankylosing spondylitis: implications for treatment outcomes. *Arthritis Rheum.* 2008;58:649–56.
19. Wanders A, Heijde D, van der, Landewé R, Béhier J-M, Calin A, et al. Non-steroidal antiinflammatory drugs reduce radiographic progression in patients with ankylosing spondylitis: a randomized clinical trial. *Arthritis Rheum.* 2005;52:1756–65.