

Walter Alberto Sifuentes-Giraldo,^{a,*} Cristina Clara Macía-Villa,^b
Mónica Vázquez-Díaz^a

^a Servicio de Reumatología, Hospital Universitario Ramón y Cajal, Madrid, España

^b Servicio de Reumatología, Hospital Clínico Universitario de Valladolid, Valladolid, España

* Corresponding author.

E-mail address: albertosifuentesg@gmail.com
(W.A. Sifuentes-Giraldo).

<http://dx.doi.org/10.1016/j.reuma.2016.05.005>

2173-5743/

© 2016 Elsevier España, S.L.U. and Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. All rights reserved.

Similarities Between Takayasu Arteritis and Giant Cell Arteritis*



Similitudes entre la arteritis de Takayasu y la arteritis de células gigantes

Dear Editor,

We were very interested in the letter to the editor published in this journal in March of this year, by Drs. Martín Guillén, Álvarez de Cienfuegos and Hurtado García.¹

The case they mention brought us to make certain reflections. Our patient is a 59-year-old woman at the onset of the disease, with occipital headaches accompanied by an increase in levels of acute-phase reactants. Physical examination revealed asymmetric peripheral pulses and the same was observed in temporal pulses.

Both Takayasu arteritis (TA) and giant cell arteritis (GCA) are vasculitides that predominantly affect large-sized vessels.²

There are significant differences between these 2 disorders. Takayasu arteritis is generally found in patients under 40 years of age. It mostly involves the aorta and its major branches, whereas GCA mainly affects patients over 50 years of age, and is detected in branches of the external carotid artery. Both diseases predominantly affect women, they share pathogenic mechanisms and pathological findings, and have a similar response to treatment.

The typical symptoms of GCA are headache, loss of vision, jaw claudication and symptoms of polymyalgia. However, approximately 40% of the patients have manifestations considered atypical. Within this group, clinical involvement of the aorta and its major branches is found in 10%–15% of the patients as an initial manifestation.³ A study performed by Ostberg in autopsies of patients with GCA revealed the involvement of the aorta in 12 of 13 cases.⁴ In their cohort of 168 patients with GCA, Nueninghoff et al. found that 27% had complications affecting the large vessels. In all, 18% had aortic aneurysm/dissection and 13% had stenosis.⁵ The majority of the cases are asymptomatic and it is underdiagnosed and, thus, its true prevalence is unknown. Patients with stenosis of the large vessels generally have fewer cranial symptoms, are found to have fewer changes in the temporal arteries on biopsies and a less marked increase in the levels of acute-phase reactants.⁵

The detection of the involvement of the aorta and its branches increases notably when ancillary tests like ultrasound, computed tomography angiography, positron emission tomography with ¹⁸F-fluorodeoxyglucose (FDG-PET) or angiography are utilized.^{3,6–8}

The study evaluated 35 GCA patients who had not been treated with corticosteroids because of FDG-PET and found that 83% had increased FDG uptake in the large arteries. The involvement of the subclavian artery and aorta was observed in 74% and 54% of

the cases, respectively. The uptake of FDG had increased after 3 months of treatment with corticosteroids.⁶ In another study, computed tomography angiography showed involvement of the large vessels in 67.5% of 40 patients with GCA. Those most widely affected were the aorta (65%), brachiocephalic trunk (47.5%), carotid arteries (35%) and subclavian arteries (42.5%).⁷

In short, their patient, at the age of 64 years, has headaches but no Doppler evidence or temporal artery biopsy, and the predominating signs are arterial stenoses. She may have TA, as was suggested, or could have GCA, with stenoses involving the large vessels.

References

- Martín Guillén S, Álvarez de Cienfuegos A, Hurtado García R. Nuevos fármacos en la arteritis de Takayasu, papel del tocilizumab. *Reumatol Clin.* 2016; <http://dx.doi.org/10.1016/j.reuma.2016.01.008>.
- Jennette JC, Falk RJ, Bacon PA, Basu N, Cid MC, Ferrario F, et al. 2012 revised International Chapel Hill Consensus Conference nomenclature of vasculitides. *Arthritis Rheum.* 2013;65:1–11.
- Cid MC, Prieto-González S, Arguis P, Espigol-Frigolé G, Butjosa M, Hernández-Rodríguez J, et al. The spectrum of vascular involvement in giant-cell arteritis: clinical consequences of detrimental vascular remodeling at different sites. *APMIS.* 2009; Suppl. 127:10–20.
- Ostberg G. Morphological changes in the large arteries in polymyalgia arteritica. *Acta Med Scand Suppl.* 1972;533:135–59.
- Nueninghoff DM, Hunder GG, Christianson TJ, McClelland RL, Matteson EL. Incidence and predictors of large-artery complication (aortic aneurysm, aortic dissection, and/or large-artery stenosis) in patients with giant cell arteritis: a population-based study over 50 years. *Arthritis Rheum.* 2003;48:3522–31.
- Blockmans D, de Ceuninck L, Vanderschueren S, Knockaert D, Mortelmans L, Bobbaers H. Repetitive ¹⁸F-fluorodeoxyglucose positron emission tomography in giant cell arteritis: a prospective study of 35 patients. *Arthritis Rheum.* 2006;55:131–7.
- Prieto-González S, Arguis P, García-Martínez A, Espigol-Frigolé G, Tavera-Bahillo I, Butjosa M, et al. Large vessel involvement in biopsy-proven giant cell arteritis: prospective study in 40 newly diagnosed patients using CT angiography. *Ann Rheum Dis.* 2012;71:1170–6.
- Schmidt WA, Seifert A, Gromnica-Ihle E, Krause A, Natusch A. Ultrasound of proximal upper extremity arteries to increase the diagnostic yield in large-vessel giant cell arteritis. *Rheumatology (Oxford).* 2008;47:96–101.

Maria Noelia Antoniol, Carlos Edgardo Perandones*

Sección Reumatología, Departamento de Medicina Interna, Fundación para la Lucha contra las Enfermedades Neurológicas de la Infancia (FLENI), Buenos Aires, Argentina

* Corresponding author.

E-mail address: perandones0@gmail.com (C.E. Perandones).

<http://dx.doi.org/10.1016/j.reuma.2016.05.004>

2173-5743/

© 2016 Elsevier España, S.L.U. and Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. All rights reserved.

* Please cite this article as: Antoniol MN, Perandones CE. Similitudes entre la arteritis de Takayasu y la arteritis de células gigantes. *Reumatol Clin.* 2017;13:122.