




Sociedad Española
de Reumatología -
Colegio Mexicano
de Reumatología

Reumatología Clínica

www.reumatologiaclinica.org



Original Article

CT-Guided Needle Biopsy in Vertebral Osteomyelitis: Study of Factors That Could Influence in Culture Yield[☆]



Jorge Juan Fragío Gil^{a,b,*}, Roxana González Mazarío^a, José Ivorra Cortés^a,
Antonio José Cañada Martínez^b, Miguel Salavert Lleti^c, José Andrés Román Ivorra^a

^a Rheumatology Department, Hospital Universitari i Politècnic La Fe, Valencia, Spain

^b Instituto de Investigación Sanitaria La Fe (IIS La Fe), Valencia, Spain

^c Infectious Diseases Unit, Hospital Universitari i Politècnic La Fe, Valencia, Spain

ARTICLE INFO

Article history:

Received 25 May 2020

Accepted 17 August 2020

Keywords:

Vertebral osteomyelitis
CT guided biopsy
Spondylodiscitis

ABSTRACT

Objective: The aim of this study was to analyze which are the main factors that could influence the result of a CT guided biopsy in vertebral osteomyelitis (VO) patients.

Methods: A single center retrospective observational study was performed including adult patients who had been diagnosed with VO and undergone CT guided needle biopsy from January 2010 to January 2020. Demographical features, concurrent diseases, laboratory findings, microbiological diagnosis, radiological data, medical complications, antibiotic exposure were compiled. Multivariate analysis was performed with a logistic regression comparing the patients depending on the culture result.

Results: Seventy-seven patients were included in the study. Baseline characteristics were comparable between groups. Sample culture was positive in 43 cases (56%). Microorganism isolated were gram + (72%), gram – (14%), mycobacteria (7%) and fungi (7%). Delay in the procedure, antibiotic exposure and blood culture positivity were also similar among both groups. The biopsy results were not influenced by the CRP value, the presence of fever nor antibiotic exposure. The longer duration of back pain was associated to a lower probability of a positive culture.

Conclusions: In conclusion, our study displays an acceptable reliability of CT guided needle biopsy in VO patients, even in cases under antibiotic treatment. The presence of fever or CRP values did not predict a positive culture. Delay in diagnosis could impact negatively on culture yield.

© 2020 Elsevier España, S.L.U. and Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. All rights reserved.

Punción asistida por TC en pacientes con osteomielitis vertebral: análisis de los factores que condicionan su rentabilidad

RESUMEN

Objetivo: El propósito del presente estudio es analizar qué factores pueden influir en el resultado del cultivo de las muestras obtenidas por punción guiada por TC en pacientes con osteomielitis vertebral.

Métodos: Se realizó un estudio en un único centro, retrospectivo y observacional en pacientes diagnosticados de osteomielitis vertebral, que fueron subsidiarios de punción-biopsia entre enero de 2010 y enero de 2020. Se recogieron para su análisis, variables demográficas, comorbilidades, resultados de laboratorio, radiología, el tratamiento previo con antibióticos y la demora previa a la realización de la técnica. Se realizó un análisis multivariante mediante regresión logística.

Palabras clave:

Osteomielitis vertebral
Punción guiada por TC
Espindilodiscitis

[☆] All authors have seen and approved the manuscript, contributed significantly to the work, and also that the manuscript has not been previously published nor is not being considered for publication elsewhere.

* Corresponding author.

E-mail address: jorgefragio@gmail.com (J.J. Fragío Gil).

Resultados: Se incluyó a un total de 77 pacientes que fueron sometidos a la técnica. Sus características basales fueron similares. El cultivo fue positivo en 43 casos (56%). Los microorganismos aislados fueron gram + (72%), gram – (14%), micobacterias (7%) y hongos (7%). El retraso en la ejecución de la técnica y el tratamiento previo con antibióticos fue similar en ambos grupos. Ni el valor de PCR, la presencia de fiebre ni la antibioterapia tuvieron influencia en el resultado del cultivo. Se observó que una mayor duración del dolor lumbar se relacionó con una menor probabilidad de obtener un resultado positivo en el cultivo.

Conclusiones: Incluso bajo exposición antibiótica, la punción asistida por TC mostró una rentabilidad aceptable. La presencia de fiebre o valores elevados de PCR no fueron predictivos de positividad del cultivo. El retraso diagnóstico sí podría impactar negativamente en la rentabilidad diagnóstica del cultivo procedente de la biopsia.

© 2020 Elsevier España, S.L.U.

y Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. Todos los derechos reservados.

Introduction

Vertebral Osteomyelitis (VO) is an infectious disease of the vertebral body which can involve the intervertebral disc space.¹ The clinical manifestations, laboratory markers and radiological findings may be sometimes unspecific, thus its diagnosis turns frequently into a challenge.^{2,3} Therefore, current ISDA (Infectious Diseases Society of America) guidelines suggest that a proper evaluation of a patient should include a thorough neurological examination, baseline CRP and ESR determination, a spine imaging technique (preferred MRI) and bacterial blood cultures. Moreover it is recommended, if possible, to withhold the antibiotic therapy until a microbiological diagnosis could be established. However, if blood cultures do not yield the causative microorganism, an image-guided aspiration biopsy is encouraged.³ The most frequent microorganism involved is *Staphylococcus aureus* in more than half of the cases but other causes could be seen such as other bacteria species or even fungi or mycobacteria.^{4,5}

Biopsy material could be obtained via open procedure (open biopsy) or by percutaneous access (usually assisted by CT or fluoroscopy). There are several studies reporting both techniques yield, which had shown a wide range of results. Some of them had shown low positivity rates with CT guided biopsy (30–50%) and higher with open biopsy,^{6–8} however other groups highlighted the importance of withholding antibiotics prior to the procedure,^{9–12} fact that needs to be further explored because other studies have failed in finding any relationship between the antibiotic exposure and the technique result. Regardless, though open biopsy seems to be more reliable, percutaneous procedures such as CT guided biopsy are preferred¹ because they are safer, less invasive, require shorter hospital stays and in most cases, general anesthesia is not required.¹³

Since there is still controversy around this topic,⁶ the aim of this study was to analyze which are the main factors that could influence in the result of a CT guided biopsy in vertebral osteomyelitis patients.

Patients and methods

Study design and population

A single center retrospective observational study was performed. Inclusion criteria was adult patients who had been diagnosed with VO from January 2010 to January 2020. Diagnosis of VO, according to other studies was established based on the combination of the clinical presentation with either a definitive bacteriologic diagnosis or pathological and/or imaging studies.^{2–12} Patients were admitted to different wards, so medical and surgical approach was not standardized. Patients who did not undergo CT guided biopsy or had a postsurgical infection were excluded. Coagulase-negative staphylococcus was considered

a true pathogen when it was isolated from biopsy or when it was detected in two or more blood cultures.

Clinical cases were search through the clinical history management software (Orion Clinic, Everis Group). Terms searched were “vertebral osteomyelitis”, “discitis” and “spondylodiscitis”.

Demographical features, concurrent diseases, laboratory findings, microbiological diagnosis, radiological data, medical complications, antibiotic exposure were compiled.

Variable definition

Immunosuppressed patients were considered those who had a rheumatic or inflammatory bowel disease undertaking immunomodulatory drugs, solid organ transplantation receptors, or in Human Immunodeficiency Virus (HIV) patients. Neurological deficit included paresthesia, limb weakness, radiculopathy, sensory loss and/or paraparesis. Clinical and radiological history of lumbar stenosis or disc herniation was considered as prior spine pathology. Chronic Kidney Disease was considered when glomerular filtration rate (GFR) was below 30 mL/min, thus, pertaining to groups 4 and 5 of the CKD classification.

Radiological presence of epidural abscess, paravertebral abscess, cord compression and vertebral destruction was determined by a musculoskeletal specialized radiologist. Infection location was defined by the most superior segment involved.

Procedure description

Standard procedure at our center is performed by Musculoskeletal Specialized Radiologist under local anesthesia and CT control. Abscess sample is collected with Acecut® needle (18G) with coaxial technique, trying to obtain at least 3 samples. For discal space, Ostycut® needle (13.5G–15G) is used. For detection of mycobacteria, culture and real time PCR was performed (Seegene kit). If possible, samples were sent for histopathology, anaerobic, aerobic, mycobacteria and fungal cultures.⁵

Statistical analysis

Continuous variables were summarized using mean (SD, standard deviation) and median (1st, 3rd quartiles). Categorical variables were summarized using absolute and relative frequencies (%). To assess the association between risk factors a logistic regression was adjusted including CRP at admission, the presence of fever, back pain span and antibiotic exposure comparing the patients depending on the culture result. All statistical analyses were performed using software R (version 3.5.0). *P* values were set based on number of analyses.




Fig. 1. General study procedure. From a total of 122 cases of VO at our centre, 45 were excluded and 77 included in the study. In 43 patients the CT guided needle biopsy yield the microbiological diagnosis.

Results

Study patients

Overall, from a total of 122 vertebral osteomyelitis cases found, 77 (63%) were included in the study (Fig. 1). Patients were divided into groups depending on the culture result. Baseline characteristics are shown below (Table 1).

Most patients were diagnosed with MRI (65% and 85% respectively between groups), followed by CT (33% and 15% respectively). In 1 case in the biopsy positive group, 18F-FDG PET/CT was the diagnosis imaging tool.

Intervention

Delay in the procedure, antibiotic exposure and blood culture positivity were similar among both groups (Table 2).

Sample culture was positive in 43 cases (56%) as seen in Fig. 1. Microorganism isolated included gram + (72%), gram – (14%), mycobacteria (7%) and fungi (7%). Detailed data of microbiological findings are provided in Table 3. Seventeen patients had a positive result on the biopsy and blood cultures. In 14 of these 17 cases (83%) it was isolated the same organism on the blood and biopsy culture (see notations below Table 3). Microbiological diagnosis in biopsy negative group was established according to blood cultures and was possible in 16 patients (47% of the biopsy negative group).

Multivariate analysis was performed with a logistic regression including CRP at admission, fever, back pain span and recent antibiotic treatment. The biopsy cultures were not influenced by the CRP value (CI 95% [0.994–1.006] *p*=0.927), the presence of fever (CI 95% [0.518–5.533] *p*=0.41) nor antibiotic exposure (CI 95% [0.116–1.355] *p*=0.155). The longer duration of the back pain was associated to a lower probability of a positive biopsy (CI 95% [0.977–0.998] *p*=0.042).

Patients received antibiotic treatment during the next 84 (56, 112) days in biopsy positive group and 84 (56, 120) days in negative group. Fifteen patients required surgery (6 in positive group and 9 in the negative group, 14% and 26% respectively). Sixteen patients were readmitted after discharge (16 in positive group and 10 in the negative group, 37% and 29% respectively). Six patients died due

Table 1
Demographic and baseline characteristics.

Characteristic	Biopsy culture positive (56%) n = 43	Biopsy culture negative (44%) n = 34
Age – years (ds)	60 (16.6)	61 (14.2)
Male sex – number (%)	33 (77)	21 (62)
Comorbidities – number (%)		
High blood pressure	21 (49)	18 (59)
Diabetes Mellitus	12 (28)	5 (15)
Chronic kidney disease	7 (16)	7 (21)
Liver cirrhosis	7 (16)	8 (24)
Immunosuppression – number (%)	9 (21)	8 (24)
Prior spine pathology – number (%)	20 (47)	19 (56)
Symptoms and signs at diagnosis		
Back pain – number (%)	43 (100)	33 (97)
Time since pain started – median	30 (20.5, 60)	55 (22.5, 90)
Fever – number (%)	15 (38)	12 (36)
Time since fever started – median	0 (0, 1)	0 (0, 6)
Acute paraparesis – number (%)	5 (12)	7 (21)
Underlying endocarditis – number (%)	0 (0)	2 (6)
Acute phase reactants		
CRP – mg/L median	88.8 (48.9, 124)	73.8 (23.1, 128)
ESR – mm/h mean	69.5 (32.1)	59.5 (24.3)
Infection location – number (%)		
Cervical	5 (12)	4 (12)
Dorsal	19 (44)	14 (41)
Lumbo-sacral	19 (44)	16 (47)
Abscess – number (%)	42 (98)	32 (94)
Paravertebral	41 (95)	32 (94)
Epidural	29 (67)	21 (62)
Vertebral destruction – number (%)	29 (67)	19 (56)
Cord compression – number (%)	15 (35)	17 (50)

Mean values are expressed: Mean (SD)/n(%). Median values are expressed: Median (1st, 3rd Q.).

Table 2
Procedure related variables.

	Biopsy culture positive (56%) n = 43	Biopsy culture negative (44%) n = 34
Blood culture (positive) – number (%)	14 (42%)	13 (41%)
Time since admission to procedure – median days	6 (3, 11)	6.5 (3, 18.5)
Antibiotic exposure – number (%)	19 (49%)	18 (58%)
Antibiotic exposure prior procedure – days	0 (0, 6)	4 (0, 11)

to the infection (4 cases) or related complications (2 cases) (3 in positive group and 3 in the negative group, 7% and 9% respectively).

Immunosuppressed patients

A total of 17 patients were immunosuppressed. Results of these population are commented separately due to its special characteristics. Immunosuppressed patients represented the 22% of the total sample and were balanced between both groups, whom exhibited similar baseline characteristics than non-immunosuppressed patients (see Table 1). Acute phase markers seemed to be slightly inferior in negative biopsy group. Positive biopsy group may have more abscess formation (100% of the positive group and 87% on the negative group, but epidural abscess was detected in 100% of the patients of the positive group and in the 50% of the negative group). Delay in procedure performance was slightly inferior in immunosuppressed patients (mean 5.2 days). Gram positive predominated on cultures (8 cases) and in the other case, *C. auris*

Table 3
Microorganisms isolated from biopsies.

Microorganism isolated	n = 43
Gram positive species – number (%)	31 (72)
<i>Staphylococcus aureus</i> MS	7
<i>Staphylococcus epidermidis</i> ^a	12
<i>Staphylococcus hominis</i>	1
<i>Streptococcus mitis</i>	1
<i>Streptococcus anginosus</i>	2
<i>Streptococcus beta hemol grupo B</i>	1
<i>Streptococcus gallolyticus</i> spp. ^b	1
<i>Streptococcus sanguinis</i> ^c	1
<i>Corynebacterium</i> spp.	1
<i>Staphylococcus capitis</i> ^d	2
<i>Enterococcus</i> spp.	1
<i>Propionibacterium acnes</i>	1
Gram negative species – number (%)	6 (14)
<i>Pseudomona aeruginosa</i> ^e	3
<i>Brevundimonas</i> spp.	1
<i>Escherichia coli</i> ^f	2
Fungi – number (%)	3 (7)
<i>Candida auris</i> ^g	2
<i>Candida albicans</i> ^h	1
Mycobacteria – number (%)	3 (7)
<i>Micobacterium tuberculosis</i>	3

^a In 4 cases were also detected on the blood cultures.

^b Also was isolated on the blood cultures of a patient who was also diagnosed of colorectal cancer.

^c Also was isolated on the blood cultures of a patient.

^d Also was isolated on the blood cultures of a patient.

^e One of them was a multiresistant *p. Aeruginosa* that also was isolated from blood cultures.

^f On the same culture of a single patient, grown *E. coli* and *Dermatobacter hominis*, *E. coli* was considered the true pathogen. On the other case, *E. coli* was also isolated on the blood culture.

^g Also was isolated on the blood cultures of a patient.

^h Also was isolated on the blood cultures of a patient.

was detected. In 4 cases (44%), blood and biopsy cultures isolated the same microorganism. Two patients were splenectomized and in one case *Staphylococcus epidermidis* was isolated on the blood culture (during a febrile episode) and in the other no potential causative organism was detected (the biopsy was performed under antibiotic exposure). On both cases, the biopsy was negative with a delay of 3 and 4 days respectively.

Discussion

In this study including VO patients, the focus has been placed on the biopsy that is needed in many patients, in an attempt of elucidating the causative microorganism so as to provide a targeted antibiotherapy, and thus, reducing side effects and improving the efficacy.^{1,7} In particular CT guided needle biopsy had been analyzed in a population with demographic and clinical comorbidities distribution similar to other studies.^{2,6,7} The population of the study had been split in two groups attending to the result of the culture of the sample obtained from de biopsy.

In this cohort a higher presence of immunosuppressed patients could be noted, due to the characteristics of the center, which is a university hospital. The results of that subset of patients have been displayed separately. In general, the distribution of cardiovascular risk factors and other clinical variables has been fair between both groups. Although there are some differences on the radiological variables (more prevalence of epidural abscess and cord compression) and also, the acute phase markers were slightly greater on the positive result group. In any case, extravertebral extension could not act as a factor of poor prognosis.⁴ This fact would be explored on further studies but seems to be logical due to a

presumed more aggressiveness of the infection on immunocompromised host. Moreover, other studies found correlation between higher CRP value and greater chance of a positive result.^{7–9}

The signs and symptoms collected seems to be in agreement with other series^{3,4,7} with the expected high prevalence of back pain, presented in almost all patients and fever in just 1 out 3 patients, approximately in both groups. As seen in other works, delay in diagnosis is still a problem in VO patients^{2,4} who usually waits at least a month since the beginning of the pain, until diagnosis. Additionally, in some cases this period is increased dramatically. Further strategies should be set up in order to reduce this term and prevent future complications.¹⁴

Infectious endocarditis is a frequent association with VO,^{15,16} but in this study were scarcely represented because in those patients the causative microorganism is more frequent detected through the blood cultures.

Acute phase markers (CRP and ESR) were elevated in both groups. Numerically, values were higher in biopsy positive groups, as seen in other studies^{7–9} but statistical signification was not reach. Radiological variables were comparable in both groups. A slight difference was noted in cord compression (higher in negative group) probably related to a greater presence of patients with acute paraparesis at diagnosis in this group. This result might be linked to the higher presence of further surgical approach detected in the negative culture group.

As it has been pointed out in several studies delay in diagnosis is probably the main issue in VO.^{2,17} Moreover, since antibiotic withholding is widely recommended,¹ these biopsies should be performed as soon as possible. In this study a median delay of 6 days was detected on both groups, with few differences in the interquartile range (Table 2). Marschall et al. achieved a 65% of positivity with a median delay of 3 days.¹⁰ Better mean values in immunosuppressed patients have been noted, but overall, this should be improved and analyzed (for instance, creating a specific circuit).

Microorganisms isolated had a distribution comparable to another studies.^{2,18} Antibiotic exposure was registered in half of the cases on the positive group and in the 58% of the negative group. The higher prevalence of patients with acute paraparesis at diagnosis could be a reasonable cause. In addition, more antibiotic exposure is presumed to reduce the culture yield,¹¹ but it was not detected on our study. This is a matter of controversy, but there are other several studies that concluded the same.^{12,17} A possible explanation of the lack or the small influence of antibiotic exposure on the culture result could be that intervertebral space is avascular. A high correlation between blood cultures and biopsy had been noted, so as it is had been previously recommended on the guidelines and other works,¹ CT-guided biopsy could be avoided in most cases.

Overall positivity (56%) was on the average of the studies published which vary from 21% to 89%.^{4,7–9,19–21} The biopsy result was not modified by prior CRP values, fever as clinical presentation nor antibiotic exposure. The longer the length of painful period before diagnosis was related to a lower chance of obtaining a positive result on culture. This result could be explained by a greater aggressiveness of pyogenic organisms that perhaps congregate in the lesser time span instead of non-pyogenic agents and seems logical that the delay in diagnosis makes a negative impact on the technique reliability. Other studies has been performed in order to detect predictive factors of positivity, such as a recent study found that using larger diameter biopsy needle increases the likelihood of causative microorganism.¹⁹ Other factor detected related to higher chance of positivity has been the attenuation value on CT scan.²⁰

In order to point out the limitations of the study, since this is a retrospective study, clinical cases may be missed during the search process. Moreover, patients were admitted to different wards, and that hence on variability in patient management. Concerning the procedure, adding more variables would provide more informa-

tion such as areas specifically punctured, specific antimicrobial treatment received prior to the procedure or number of samples taken at each procedure. During statistical analysis some difficulties were found, such a huge variability in days of antimicrobial therapy received prior the procedure, finally solved dichotomizing in “exposure: yes/no” and reducing the statistical power. The inclusion of mycobacteria and fungi might influence the clinical picture and the yield of the CT-guided biopsy.

In conclusion our study displays an acceptable reliability of CT guided needle biopsy in VO patients, even in cases under antibiotic treatment. The presence of fever or elevated CRP values did not predict a positive culture in our study. A positive blood culture would provide a microbiological diagnosis in most cases. Delay in diagnosis could impact negatively in culture yield. Therefore, improving diagnostic strategies in general and shortening the span from VO diagnosis to the procedure performance in particular, may enhance not only the culture yield reliability but also the patient prognosis.

Funding

None funding source was required.

Conflict of interest

None.

Acknowledgements

To our partners from the Rheumatology department, the Infectious disease Unit and the Musculoskeletal Unit of the Radiology department. Also to Mrs. Paula Giménez for her effort and valuable help.

The authors are grateful for the collaboration with English language revision to Mr. Paul Godfrey.

References

1. Berbari EF, Kanj SS, Kowalski TJ, Darouchi RO, Widmer AF, Schmitt SK, et al. 2015 Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines for the Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults. *Clin Infect Dis*. 2015;61:e26–46, <http://dx.doi.org/10.1093/cid/civ482>.
2. Fragió Gil JJ, González Mazarío R, Salavert Lleti M, Román Ivorra JA. Vertebral osteomyelitis: clinical, microbiological and radiological characteristics of 116 patients [published online ahead of print, 2020 May 20]. *Med Clin (Barc)*. 2020, <http://dx.doi.org/10.1016/j.medcli.2019.12.029>. S0025-7753(20)30187-1.
3. Mylona E, Samarkos M, Kakalou E, Fanourgiakis P, Skoutelis A. Pyogenic vertebral osteomyelitis: a systematic review of clinical characteristics. *Semin Arthritis Rheum*. 2009;39:10–7, <http://dx.doi.org/10.1016/j.semarthrit.2008.03.002>.
4. Nolla JM, Ariza J, Gómez-Vaquero C, Fiter J, Bermejo J, Valverde J, et al. Spontaneous pyogenic vertebral osteomyelitis in nondrug users. *Semin Arthritis Rheum*. 2002;31:271–8, <http://dx.doi.org/10.1053/sarh.2002.29492>.
5. Lew DP, Waldvogel FA. Osteomyelitis. *Lancet*. 2004;364:369–79, [http://dx.doi.org/10.1016/S0140-6736\(04\)16727-5](http://dx.doi.org/10.1016/S0140-6736(04)16727-5).
6. Nickerson EK, Sinha R. Vertebral osteomyelitis in adults: an update. *Br Med Bull*. 2016;117:121–38, <http://dx.doi.org/10.1093/bmb/ldw003>.
7. Chong BSW, Brereton CJ, Gordon A, Davis JS. Epidemiology microbiological diagnosis, and clinical outcomes in pyogenic vertebral osteomyelitis: a 10-year retrospective cohort study. *Open Forum Infect Dis*. 2018;5:ofy037, <http://dx.doi.org/10.1093/ofid/ofy037>. Published 2018 Mar 10.
8. Ang MT, Wong GR, Wong DR, Clements W, Joseph T. Diagnostic yield of computed tomography-guided biopsy and aspiration for vertebral osteomyelitis. *J Med Imaging Radiat Oncol*. 2019;63:589–95, <http://dx.doi.org/10.1111/1754-9485.12923>.
9. Marshall J, Bhavan KP, Olsen MA, Fraser VJ, Wright NM, Warren DK. The impact of prebiopsy antibiotics on pathogen recovery in hematogenous vertebral osteomyelitis. *Clin Infect Dis*. 2011;52:867–72, <http://dx.doi.org/10.1093/cid/cir062>.
10. Lee HT, Pukenas BA, Sebros R. Change in bone CT attenuation and C-reactive protein are predictors of bone biopsy culture positivity in patients with vertebral discitis/osteomyelitis [published online ahead of print, 2020 Mar 19]. *Spine (Phila Pa 1976)*. 2020, <http://dx.doi.org/10.1097/BRS.0000000000003504>, 10.1097.
11. Kim CJ, Song KH, Park WB, Suk Kim E, Won Park S, Kim HB, et al. Microbiologically and clinically diagnosed vertebral osteomyelitis: impact of prior antibiotic exposure. *Antimicrob Agents Chemother*. 2012;56:2122–4, <http://dx.doi.org/10.1128/AAC.05953-11>.
12. Crisolago PA, La Fontaine J, Wukich DK, Kim PJ, Oz OK, Lavery LA. The effect of withholding antibiotics prior to bone biopsy in patients with suspected osteomyelitis: a meta-analysis of the literature. *Wounds*. 2019;31:205–12.
13. McNamara AL, Dickerson EC, Gomez-Hassan DM, Cinti SK, Srinivasan A. Yield of image-guided needle biopsy for infectious discitis: a systematic review and meta-analysis. *AJNR Am J Neuroradiol*. 2017;38:2021–7.
14. Gupta A, Kowalski TJ, Osmon DR, Enzler M, Steckelberg JM, Huddleston PM, et al. Long-term outcome of pyogenic vertebral osteomyelitis: a cohort study of 260 patients. *Open Forum Infect Dis*. 2014;1, <http://dx.doi.org/10.1093/ofid/ofu107>, ofu107. Published 2014 Dec 5.
15. Pigrau C, Almirante B, Flores X, Falcó V, Rodríguez D, Gasser I, et al. Spontaneous pyogenic vertebral osteomyelitis and endocarditis: incidence, risk factors, and outcome. *Am J Med*. 2005;118:1287, <http://dx.doi.org/10.1016/j.amjmed.2005.02.027>.
16. Koslow M, Kuperstein R, Eshed I, Perelman M, Maor E, Sidi Y. The unique clinical features and outcome of infectious endocarditis and vertebral osteomyelitis co-infection. *Am J Med*. 2014;127, <http://dx.doi.org/10.1016/j.amjmed.2014.02.023>, 669.e9–e15.
17. Saravolatz LD 2nd, Labalo V, Fishbain J, Szpunar S, Johnson LB. Lack of effect of antibiotics on biopsy culture results in vertebral osteomyelitis. *Diagn Microbiol Infect Dis*. 2018;91:273–4, <http://dx.doi.org/10.1016/j.diagmicrobio.2018.02.017>.
18. Amsilli M, Epaulard O. How is the microbial diagnosis of bacterial vertebral osteomyelitis performed? An 11-year retrospective study [published online ahead of print, 2020 Jun 26]. *Eur J Clin Microbiol Infect Dis*. doi:10.1007/s10096-020-03929-1.
19. Hussein JS, Simeone FJ, Nelson SB, Chang CY. CT-guided discitis-osteomyelitis biopsies: needle gauge and microbiology results [published online ahead of print, 2020 Apr 18]. *Skeletal Radiol*. 2020;10, <http://dx.doi.org/10.1007/s00256-020-03439-3>.
20. Beroukhim G, Shah R, Bucknor MD. Factors predicting positive culture in CT-guided bone biopsy performed for suspected osteomyelitis. *AJR Am J Roentgenol*. 2019;212:620–4, <http://dx.doi.org/10.2214/AJR.18.20125>.
21. Sertic M, Parkes L, Mattiassi S, Pritzker K, Gardam M, Murphy K. The efficacy of computed tomography-guided percutaneous spine biopsies in determining a causative organism in cases of suspected infection: a systematic review. *Can Assoc Radiol J*. 2019;70:96–103, <http://dx.doi.org/10.1016/j.carj.2018.09.003>.