




Sociedad Española
de Reumatología -
Colegio Mexicano
de Reumatología

Reumatología Clínica

www.reumatologiaclinica.org



Images in Clinical Rheumatology

Tophaceous gout in a woman[☆]

Gota tofácea en una mujer

María José Pérez Galán,* Celia García Jiménez

Unidad de Reumatología, Complejo Hospitalario de Jaén, Jaén, Spain



ARTICLE INFO

Article history:

Received 27 May 2018

Accepted 2 August 2018

Available online 29 October 2020

Clinical case

A 76 year-old woman, with a history of anaemia and hyperuricemia treated with ferrous sulphate and 100 mg allopurinol per day. She states that she has consumed half a litre of alcohol-free beer per day for 15 years. She states that she has no history of arterial hypertension, no chronic disease and that she does not consume diuretics.

She visited our surgery due to arthralgia in the hands and feet with a crisis of pain and functional impotency in multiple joints that has evolved over more than 15 years. Physical examination found major tophaceous deposits in the hands and feet, most strikingly in the right hand (Fig. 1), with incapacity of grasping, above all in the right hand.

Analysis detected 7.7 mg/dl uricaemia, while the rest of basic biochemistry was normal. The haemogram, acute phase reagents, rheumatoid factor and anti-citrulline antibodies were normal or negative.

Bone X-ray imaging showed an important increase in the soft parts, with trapping and erosion in the metacarpophalangeal, proximal and distal interphalangeal joints of both hands (Fig. 2), as well as erosion on the first metatarsal joints and luxation of the 5th left metatarsal joint (Fig. 3).

Diagnosis

Tophaceous gout.



Figure 1. Tophaceous deposits in the hands.



Figure 2. X-ray image of hands: increased soft parts and erosion.

[☆] Please cite this article as: Pérez Galán MJ, García Jiménez C. Gota tofácea en una mujer. Reumatol Clin. 2020;16:506-507.

* Corresponding author.

E-mail address: mjg77@hotmail.com (M.J. Pérez Galán).



Figure 3. X-ray image of feet: erosion in the first metatarsophalangeal joints.

Evolution

Treatment with hygienic and dietary methods was prescribed, with colchicine and an increased dose of allopurinol to maintain uricaemia below 5 mg/dl.

Discussion

Gout arthropathy is more common in men¹ than it is in women, although it is the most frequent inflammatory disease of the joints in post-menopausal women. The first manifestation of the disease is typically acute arthritis.² Tophi are sodium urate monohydrate deposits surrounded by a granulomatous inflammatory reaction. They are a pathognomonic characteristic of gout and can be detected by physical examination and/or imaging techniques.³

We present the case of a woman with tophaceous arthropathy. We recall the importance of maintaining suitable levels of uricaemia, as this disease is potentially disabling. We underline that although it occurs more frequently in men, it is the most common inflammatory arthropathy in post-menopausal women.

Conflict of interests

The authors have no conflict of interests to declare.

References

1. Bhole V, de VM Rahman MM, Krishnan E, Choi H. Epidemiology of gout in women: Fifty-two-year followup of a prospective cohort. *Arthritis Rheum.* 2010;62:1069–76.
2. Neogi T. Clinical practice. Gout *N Engl J Med.* 2011;364:443–52.
3. Tophi, mainly in articular, periarticular, bursal, bone, auricular, and cutaneous tissues are a pathognomonic feature of gout, and are detectable by physical exam, and/or by imaging approaches and pathology examination.